


53012-32-107
ORC 68-7
APRIL 1968

A NOTE ON TESTS FOR MONOTONE FAILURE RATE BASED ON INCOMPLETE DATA

by

RICHARD E. BARLOW and FRANK PROSCHAN

AD669111



DDC
RECEIVED
MAY 22 1968
B

OPERATIONS
RESEARCH
CENTER

This document has been approved for public release and sale; its distribution is unlimited. The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

UNIVERSITY OF CALIFORNIA • BERKELEY

AD 669 111

A NOTE ON TESTS FOR MONOTONE FAILURE RATE BASED
ON INCOMPLETE DATA

Richard E. Barlow, et al

California University
Berkeley, California

April 1968

A NOTE ON TESTS FOR MONOTONE FAILURE RATE BASED ON INCOMPLETE DATA

by

Richard E. Barlow[†]
Operations Research Center
University of California, Berkeley

and

Frank Proschan
Boeing Scientific Research Laboratories
Seattle, Washington

This document has been approved for public release and sale; its distribution is unlimited. The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

April 1968

GRC 68-7

[†] This research has been partially supported by the Office of Naval Research under Contract Nonr-3656(18), the National Science Foundation under Grant GK-1684, the National Science Foundation under Grant GP-7417, and the U.S. Army Research Office-Durham, Contract DA-31-124-ARO-D-331 with the University of California. Reproduction in whole or in part is permitted for any purpose of the United States Government.

ACKNOWLEDGEMENTS

We would like to acknowledge the help of T. A. Bray,
R. Pyke, and R. Wolff.

ABSTRACT

Tests for exponential versus IFRA distributions based on incomplete data are defined and shown to be unbiased. The tests are motivated by a class of tests considered in detail by Bickel and Doksum. Tests for exponential versus IFR distributions based on the ranks of total time on test statistics are also considered.

A NOTE ON TESTS FOR MONOTONE
FAILURE RATE BASED ON INCOMPLETE DATA

by

Richard E. Barlow[†] and Frank Proschan

1. INTRODUCTION AND SUMMARY

Let $0 \equiv X_{(0)} \leq X_{(1)} \leq X_{(2)} \leq \dots \leq X_{(n)}$ be the order statistics of a (complete) random sample from a population with distribution F and density f such that $F(0) = 0$. Bickel and Doksum (1968) consider the problem of testing

$$H_0 : F(t) = 1 - e^{-\lambda t} \quad t \geq 0, \lambda > 0$$

versus

$$H_1 : F \text{ IFR}$$

(i.e., $-\log[1 - F(t)]$ convex on $[0, \infty)$).

Let $D_i = (n-i+1)(X_{(i)} - X_{(i-1)})$, $i = 1, 2, \dots, n$. They consider tests based on statistics of the form

$$\frac{\sum_{i=1}^n a_i D_i}{\sum_{i=1}^n D_i}$$

where $a_1 \geq a_2 \geq \dots \geq a_n$. The test, ϕ_a , rejects H_0 when

$$\frac{\sum_{i=1}^n a_i D_i}{\sum_{i=1}^n D_i} \geq c_{a,a,n}. \quad \text{They compute the asymptotic relative efficiency of}$$

[†]Research partially supported by the Office of Naval Research, Contract Nonr-3656(18) with the University of California, and the Boeing Scientific Research Laboratories. This report is also appearing as a Boeing Scientific Research Laboratories Document.

various such tests relative to selected parametric alternatives. Such tests were shown to be unbiased against IFRA (for increasing failure rate average) alternatives by Barlow and Proschan (1966) and hence a fortiori for IFR alternatives. [See also Birnbaum, Esary and Marshall (1965) for justification of the IFRA assumption.]

The purpose of this note is to show that analogous tests designed to treat incomplete samples of failure data are also unbiased against IFRA alternatives. Let X_i be the time to failure of the i^{th} item in a sample of size n . Let L_i be a given truncation time for the i^{th} item and let

$$Z_i = \min(X_i, L_i) \quad i = 1, 2, \dots, n.$$

Let $0 \equiv Z_{(0)} \leq Z_{(1)} \leq \dots \leq Z_{(k)}$ be the first k observed failure times. Note that "withdrawals" may occur between $Z_{(i)}$ and $Z_{(i+1)}$ and that k is, in general, a random variable. Let $n(u)$ be the (random) number of items on test at time u .

We define a test, ϕ_a^* , (a modification of ϕ_a) which rejects H_0 in favor of

$$H_1' : F \text{ IFRA}$$

(i.e., $-\{\log[1 - F(t)]\}/t$ nondecreasing on $[0, \infty)$) when

$$W_a = \frac{\sum_{i=1}^k a_i \int_{Z_{(i-1)}}^{Z_{(i)}} n(u) du}{\int_0^{Z_{(k)}} n(u) du} \geq c_{a,a,k}^*.$$

Note that $\int_{Z_{(i-1)}}^{Z_{(i)}} n(u) du$ represents the total time on test between the $(i-1)^{\text{st}}$ and

i^{th} observed failures. The distribution of W_a can be computed under H_0 using

the fact that $Y_i = \int_{Z_{(i-1)}}^{(Z_{(i)})} n(u)du$ ($i = 1, 2, \dots, k$) are distributed as

independent exponential random variables under H_0 conditioned on the value of k .

We show that ϕ_a^* is an unbiased test for IFRA alternatives for weights

$a = (a_1, a_2, \dots, a_n)$ for which $a_1 \geq a_2 \geq \dots \geq a_n$.

2. DISTRIBUTION OF W_a UNDER H_0

Let $r(t) = f(t)/[1 - F(t)]$ be the failure rate function for F . We will need the following lemma, stated without proof in Bray, Crawford, and Proschan (1967).

Lemma 1.

For any distribution F ($F(0) = 0$) with failure rate $r(t)$,

$$Y_i = \int_{Z_{(i-1)}}^{Z_{(i)}} r(u)n(u)du, \quad i = 1, 2, \dots, k \text{ are independently distributed with}$$

density e^{-y} .

Proof:

$$\text{Let } Y_1 = \int_0^{Z_{(1)}} r(u)n(u)du \text{ and } S_0(t) = \int_0^t r(u)n(u)du. \text{ Note that } S_0(t)$$

is well defined up to the time of the first observed failure since $n(u)$ depends only on the specified truncation times L_i ($i = 1, 2, \dots, n$) up until $Z_{(1)}$.

Then

$$\begin{aligned} P[Y_1 > y_1] &= P[S_0(Z_{(1)}) > y_1] = P[Z_{(1)} > S_0^{-1}(y_1)] = \\ &= \exp[-S_0(S_0^{-1}(y_1))] = e^{-y_1}. \end{aligned}$$

Thus Y_1 has density e^{-y_1} .

$$\text{Now let } Y_2 = \int_{Z_{(1)}}^{Z_{(2)}} r(u)n(u)du \text{ and } S_{x_1}(t) = \int_{x_1}^t r(u)n(u)du. \text{ Note that}$$

conditionally on $Z_{(1)} = x_1$, S_{x_1} is well defined for $x_1 \leq t < Z_{(2)}$. Hence

$$\begin{aligned}
 P[Y_2 > y_2 \mid Z_{(1)} = x_1] &= P[S_{x_1}(Z_{(2)}) > y_2 \mid Z_{(1)} = x_1] = \\
 &= P[Z_{(2)} > S_{x_1}^{-1}(y_2) \mid Z_{(1)} = x_1] = \exp\left[-S_{x_1}\left(S_{x_1}^{-1}(y_2)\right)\right] = e^{-y_2}.
 \end{aligned}$$

Thus Y_2 is independent of Y_1 and also exponentially distributed with mean 1. If we continue in this manner, conditioning on previous events, we establish the lemma. ||

Under H_0 , $r(t) \equiv \lambda$ and we see from the lemma that, given k observed failures,

$$W_a \stackrel{\text{st}}{=} \frac{\sum_{i=1}^k a_i Y_i}{\sum_{j=1}^k Y_j},$$

where $\stackrel{\text{st}}{=}$ denotes stochastic equality and Y_1, Y_2, \dots, Y_k are independent, exponentially distributed random variables with unit mean.

3. UNBIASEDNESS UNDER IFRA ALTERNATIVES

We need the following lemma to establish unbiasedness. Define

$$R(t) = \int_0^t r(u)du \quad \text{and} \quad T(t) = \int_0^t n(u)du .$$

Lemma 2.

If $\frac{R(t)}{t}$ is nondecreasing in $t \geq 0$, $n(t) \geq 0$, and $\frac{T(t)}{t}$ is nonincreasing in $t \geq 0$, then

$$(i) \quad r(t) \geq \frac{\int_0^t r(u)du}{t} \geq \frac{\int_0^t r(u)dT(u)}{T(t)}$$

$$(ii) \quad \frac{\int_0^t r(u)dT(u)}{T(t)} \quad \text{is nondecreasing in } t \geq 0 ,$$

when the indicated integrals exist.

Proof:

To show (i). The first inequality follows from differentiating $\frac{R(t)}{t}$. Since $\frac{R(t)}{t} \geq 0$ is nondecreasing in $t \geq 0$, we can approximate $R(t)$ arbitrarily closely from below by a positive linear combination of functions of the form

$$R(t) = \begin{cases} 0 & 0 \leq t < x \\ t & t \geq x \end{cases}$$

[cf. Barlow, Marshall, and Proschan (1967)]. By the Lebesgue monotone convergence theorem, we need only establish the second inequality in (ii) for functions $R(t)$ of this type. Hence for $t \geq x$,

$$\frac{\int_0^t n(u) dR(u)}{T(t)} = \frac{n(x)x + \int_x^t n(u) du}{T(t)} = 1 + \frac{xn(x) - T(x)}{T(t)}.$$

This is nondecreasing in $t \geq x$ since (a) $T(t)$ is nondecreasing in $t \geq 0$, and (b) $xn(x) - T(x) \leq 0$ since $\frac{T(x)}{x}$ is nonincreasing in $x \geq 0$.

To show (ii). Clearly

$$\frac{d}{dt} \left[\frac{\int_0^t r(u)n(u) du}{\int_0^t n(u) du} \right] \geq 0$$

if and only if

$$r(t)n(t) \int_0^t n(u) du \geq n(t) \int_0^t r(u)n(u) du$$

which follows from (i). ||

Note that if $r(t)$ is nondecreasing in $t \geq 0$, then (ii) follows for all $n(t) \geq 0$.

Lemma 2 may be used in testing for IFRA in models other than the one described in the introduction; see for example the model of Bray, Crawford, and Proschan (1967).

Theorem 1.

If F is IFRA with failure rate $r(t)$ and $Z_{(1)} \leq Z_{(2)} \leq \dots \leq Z_{(k)}$ are the observed failure times, $n(t) \geq 0$ for $t \geq 0$, and $\frac{T(t)}{t} \geq 0$ is nonincreasing in $t \geq 0$, then (conditional on k),

$$W_a = \frac{\sum_{i=1}^k a_i \int_{Z_{(i-1)}}^{Z_{(i)}} n(u) du}{\int_0^{Z_{(k)}} n(u) du} \geq \frac{\sum_{i=1}^k a_i Y_i}{\sum_{i=1}^k Y_i}$$

where $a_1 \geq a_2 \geq \dots \geq a_n$ and Y_1, Y_2, \dots, Y_k are independently distributed as exponential random variables with unit mean.

Proof:

Since $n(u) \geq 0$ and $T(t)/t$ is nonincreasing, Lemma 2 applies, yielding

$$\frac{\beta_i}{\alpha_i} = \frac{\int_0^{Z_{(i)}} r(u)n(u) du}{\int_0^{Z_{(i)}} n(u) du}$$

nondecreasing in $i = 1, 2, \dots, k$. By Lemma 1 we need only show that

$$(1) \quad \frac{\sum_{i=1}^k a_i \int_{Z_{(i-1)}}^{Z_{(i)}} n(u) du}{\int_0^{Z_{(k)}} n(u) du} \geq \frac{\sum_{i=1}^k a_i \int_{Z_{(i-1)}}^{Z_{(i)}} r(u)n(u) du}{\int_0^{Z_{(k)}} r(u)n(u) du}$$

i.e.,

$$\frac{\sum_{i=1}^k a_i (\alpha_i - \alpha_{i-1})}{\alpha_k} = \frac{\sum_{i=1}^k a_i (\beta_i - \beta_{i-1})}{\beta_k}$$

where $\alpha_0 = \beta_0 \equiv 0$. Note that

$$\sum_{i=1}^k a_i (\alpha_i - \alpha_{i-1}) = (a_1 - a_2)\alpha_1 + (a_2 - a_3)\alpha_2 + \dots + a_k \alpha_k = \sum_{i=1}^k \Delta_i \alpha_i$$

where $\Delta_i = a_i - a_{i+1} \geq 0$ for $i = 1, 2, \dots, k-1$ and $\Delta_k = a_k$. Hence

$$\frac{\beta_i}{\alpha_i} \leq \frac{\beta_k}{\alpha_k} \quad \text{implies} \quad \sum_{i=1}^k \frac{\Delta_i \alpha_i}{\alpha_k} \geq \sum_{i=1}^k \frac{\Delta_i \beta_i}{\beta_k}, \text{ which proves (1). } ||$$

4. APPLICATION OF TOTAL TIME ON TEST

Assuming an exponential distribution, the results of Bickel and Doksum (1968) may be used to establish the asymptotic normality of W_a in the incomplete data case for selected vectors $a = (a_1, \dots, a_k)$. Perhaps, the most useful test is the total time on test statistic. In the case of a complete sample of size n , this is S_1^* in the Bickel-Doksum paper, obtained by choosing $a_1 = -1/(n+1)$, after algebraic manipulation. In the case of incomplete data as described in the introduction, with k failures observed; the total time on test statistic is

$$W_{a^\circ} = \frac{\sum_{i=1}^{k-1} (k-i) \int_{Z_{(i-1)}}^{Z_{(i)}} n(u) du}{\int_0^{Z_{(k)}} n(u) du},$$

obtained by choosing $a^\circ = (k-1, k-2, \dots, 1, 0)$.

The exact distribution conditioned on the number of observed failures $k \geq 2$ is easily computed in this case. Table 1 is a short table of percentage points. Note that, under H_0

$$W_{a^\circ} \stackrel{st}{=} U_1 + U_2 + \dots + U_{k-1}$$

when U_i ($i = 1, 2, \dots, k-1$) are independent uniform random variables on $[0, 1]$. Since the distribution of W_{a° is symmetric about $\frac{k-1}{2}$, we tabulate upper percentiles only.

TABLE 1: PERCENTILES χ_α OF TOTAL TIME ON TEST STATISTIC, W_{a^0}

$k-1 \backslash \alpha$.900	.950	.975	.990	.995
2	1.553	1.684	1.776	1.859	1.900
3	2.157	2.331	2.469	2.609	2.689
4	2.753	2.953	3.120	3.300	3.411
5	3.339	3.565	3.754	3.963	4.097
6	3.917	4.166	4.376	4.610	4.762
7	4.489	4.759	4.988	5.244	5.413
8	5.056	5.346	5.592	5.869	6.053
9	5.619	5.927	6.189	6.487	6.683
10	6.178	6.504	6.781	7.097	7.307
11	6.735	7.077	7.369	7.702	7.924
12	7.289	7.647	7.953	8.302	8.535

k = number of failures observed in incomplete sample

$$P[W_{a^0} \leq \chi_\alpha] = \alpha$$

5. MONOTONE TESTS UNDER IFR ALTERNATIVES

Bickel and Doksum (1968) define a test ϕ to be monotone in the normalized spacings D_1, \dots, D_n if $\phi(D'_1, \dots, D'_n) \leq \phi(D_1, \dots, D_n)$ for all (D_1, \dots, D_n) and (D'_1, \dots, D'_n) such that for $i < j$, $D'_i \geq D'_j$ implies $D_i \geq D_j$. We show

that if D_i is replaced by $\int_{Z_{(i-1)}}^{Z_{(i)}} n(u) du$ in the incomplete data case, then a

monotone test is unbiased for testing H_0 versus H_1 when $n(u) \geq 0$ for $u \geq 0$. The test rejects H_0 for large values of ϕ .

We need

Lemma 3.

Let $r(u) \uparrow$ and $n(u) \geq 0$ for $u \geq 0$. Then for $0 \leq a < b \leq c < d$,

$$\frac{\int_a^b n(u)r(u)du}{\int_a^b n(u)du} \leq \frac{\int_c^d n(u)r(u)du}{\int_c^d n(u)du}.$$

Proof:

$$\frac{\int_a^b n(u)r(u)du}{\int_a^b n(u)du} \leq \frac{r(b) \int_a^b n(u)du}{\int_a^b n(u)du} \leq \frac{r(c) \int_c^d n(u)du}{\int_c^d n(u)du} \leq \frac{\int_c^d n(u)r(u)du}{\int_c^d n(u)du}. \quad ||$$

From Lemma 3, we immediately obtain

Theorem 2.

Let ϕ be a monotone test of H_0 versus H_1 based on a sample of incomplete data as described in the introduction. Then

$$E \left[\phi \left(\int_0^{Z^{(1)}} n(u) du, \dots, \int_{Z^{(k-1)}}^{Z^{(k)}} n(u) du \right) \mid F \text{ IFR} \right] \geq \\ \geq E[\phi(Y_1, \dots, Y_k) \mid F \text{ exponential}] ,$$

where Y_1, \dots, Y_k independent exponentially distributed random variables.

Proof:

For $i < j$,

$$\frac{\int_{Z^{(i-1)}}^{Z^{(j)}} n(u) du}{\int_{Z^{(i-1)}}^{Z^{(i)}} n(u) du} \leq \frac{\int_{Z^{(i-1)}}^{Z^{(j)}} r(u) n(u) du}{\int_{Z^{(i-1)}}^{Z^{(i)}} r(u) n(u) du} \stackrel{\text{st}}{=} \frac{Y_j}{Y_i} .$$

The inequality follows from Lemma 3; the stochastic equality follows from Lemma 1.

$$\text{Thus } \phi(Y_1, \dots, Y_k) \stackrel{\text{st}}{\leq} \phi \left(\int_0^{Z^{(1)}} n(u) du, \dots, \int_{Z^{(k-1)}}^{Z^{(k)}} n(u) du \right) . \text{ The}$$

conclusion follows by taking expectations. ||

REFERENCES

- [1] Barlow, R. E. and F. Proschan, "Inequalities for Linear Combinations of Order Statistics from Restricted Families," Ann.Math.Statist., Vol. 37, pp. 1574-1592, (1966).
- [2] Barlow, R. E., A. W. Marshall and F. Proschan, "Inequalities for Starshaped and Convex Functions," Boeing Document D1-82-0643, (1967).
- [3] Bickel, P. J. and K. Doksum, "Tests for Monotone Failure Rate Based on Normalized Spacings," Ann.Math.Statist., (to appear), (1968).
- [4] Birnbaum, Z. W., J. D. Esary and A. W. Marshall, "Stochastic Characterization of Wear-Out for Components and Systems," Ann.Math.Statist., Vol. 37, pp. 816-825, (1965).
- [5] Bray, T. A., G. B. Crawford and F. Proschan, "Maximum Likelihood Estimation of a U-Shaped Failure Rate Function," Boeing Document D1-82-0660, (1967).

Unclassified
Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1 ORIGINATING ACTIVITY (Corporate author) University of California, Berkeley		2a REPORT SECURITY CLASSIFICATION Unclassified
		2b GROUP
3 REPORT TITLE A NOTE ON TESTS FOR MONOTONE FAILURE RATE BASED ON INCOMPLETE DATA		
4 DESCRIPTIVE NOTES (Type of report and inclusive dates) Research Report		
5 AUTHOR(S) (Last name, first name, initial) BARLOW, Richard E. PROSCHAN, Frank		
6 REPORT DATE April 1968	7a TOTAL NO OF PAGES 14	7b NO OF REFS 5
8a CONTRACT OR GRANT NO. Nonr-3656(18)	9a ORIGINATOR'S REPORT NUMBER(S) ORC 68-7	
b. PROJECT NO NR 042 238		
c Research Project No.: WW 041	9b OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d		
10 AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited.		
11 SUPPLEMENTARY NOTES SEE TITLE PAGE	12 SPONSORING MILITARY ACTIVITY Mathematical Science Division	
13 ABSTRACT Tests for exponential versus IFRA distributions based on incomplete data are defined and shown to be unbiased. The tests are motivated by a class of tests considered in detail by Bickel and Doksum. Tests for exponential versus IFR distributions based on the ranks of total time on test statistics are also considered.		

DD FORM 1473
1 JAN 64

Unclassified
Security Classification

14 KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Monotone Failure Rate IFRA Distributions Exponential Total Time on Test Unbiased Tests						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.